

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Buresu



DER THE PATENT COOPERATION TREATY (PCT)

INTERNATIONAL APPLICATION PUBLISH	HED U	JNDER THE PATENT COUPERATION TREATT (101)
(51) International Patent Classification 6:		(11) International Publication Number: WO 95/33475
A61K 38/00	A1	(43) International Publication Date: 14 December 1995 (14.12.95)
(21) International Application Number: PCT/EP (22) International Filing Date: 5 June 1995 ((30) Priority Data: 9411292.7 6 June 1994 (06.06.94) (71) Applicant (for all designated States except BB US) RESEARCH & DEVELOPMENT COMPANY I [IL/IL]; P.O. Box 95, 76100 Rehovot (IL). (71) Applicant (for BB only): ORVET BV [NL/NL]; P.O. 3640, NL-AE Mijdrecht (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): KOTT, Edna Mapu, 49202 Petach Tikva (IL). KESLER, Anat [Yatkovsky, 49652 Petach Tikva (IL). (74) Agent: SCHLICH, George, William; Mathys & Sc. Grays Inn Road, London WC1X BAL (GB).	e YEL LIMITE Box 2! [IL/IL];	(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, IP, KE, KG, KP, KR, KZ, LK, LT, LU, LV, MD, MG, MN, MW, NO, YD, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NIL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.
	ATT TE A	TIPE OF A MEDICAMENT FOR THE TREATMENT OF OPTION

(54) Title: USE OF COPOLYMER-1 FOR THE MANUFACTURE OF A MEDICAMENT FOR THE TR NEURITIS

(57) Abstract

Use of copolymer-1 to treat visual impairments associated with multiple sclerosis.

Applicants: Alexander Gad and Dora Lis Serial No.: 09/816,989

Filed: March 23, 2001

Exhibit 22

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
ΑÜ	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
		GR	Greece	NL	Netherlands
BE _	Belgium	HU	Hungary	NO	Norway
BF	Burkins Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	iT	Italy	PL	Poland
BJ	Benin	JP		PT	Portugal
BR	Brazil	KE	Japan V	RO	Romania
BY	Belarus		Kenya	RU	Russian Federation
CA	Canada	KG	Kyrgystan	SD	Sudan
CF	Central African Republic	KP	-Democratic People's Republic		Sweden .
CG	Congo		of Korea	SE	•
CH	Switzerland	KR	Republic of Korea	<u> </u>	Slovenia
CI	Côte d'Ivoire	KZ	Kazakhstan	SK	Slovakia
CM	Cameroon	Li	Liechtenstein	SN	Senegal
CN	China	. LK	Sri Lanka	TD	Chad
cs	Czechoslovakia	LU	Luxembourg	TG	Togo
cz	Czech Republic	LV	Larvia	Tj	Tejikistan
DE	Germany	MC	Mosaco	TT	Trinidad and Tobago
	_ •	MD	Republic of Moldova	UA	Ukraine
DK	Denmark	MG	Madagascar	US	* United States of America
ES	Spain		Mali	. UZ	Uzbekistan
FI	Finland	ML		. VN	Viet Nam
FR	France	MN	Mongolia .	*14	V 994 1 18044
GA	Gabon				

USE OF COPOLYMER-1 FOR THE MANUFACTURE OF A MEDICAMENT.

FOR THE TREATMENT OF OPTIC NEURITIS

FIELD OF INVENTION

The present invention relates to the use of copolymer-1 in treating visual impairments associated with multiple sclerosis.

PRIOR ART

5

10

15

20

Throughout this application, various references are referred to. Disclosures of these publications in their entireties are hereby incorporated by reference into this application to more fully describe the state of the art to which this invention pertains.

Multiple sclerosis (MS) is a slow progressive CNS disease characterized by disseminated patches of demyelination in the brain and spinal cord, resulting in multiple and varied neurological symptoms and signs, usually with remissions and exacerbations (relapses). A common symptom prior to diagnosis of MS is some degree of visual impairment, frequently optic neuritis or retrobulbar optic neuritis. Alternatively, these and other visual impairments may only develop after a diagnosis of MS has been confirmed. These symptoms may also develop during the progression of the disease alongside other associated visual problems.

25

30

*****3.5

Several studies have concentrated on assessing whether optic neuritis is a reliable predictor of multiple sclerosis (Cohen MM et al., Neurology (1979) 29 208-213, Rizzo JF et al., Neurology (1988) 38 185-199, Beck RW et al., Neurology (1992) 42 1133-1135 and New Eng J Med (1993) 326 581-588).

Copolymer-1 is a synthetic polypeptide analog of myelin basic protein (MBP), which is a natural component of the myelin sheath. It has been suggested as a potential therapeutic agent for multiple sclerosis (Eur. J. Immunol. [1971] 1:242; and J. Neurol. Sci. [1977] 31:433).

10

20

25

30

35

Copolymer-1 was developed by Drs. Sela, Arnon, and their coworkers at the Weizmann Institute (Rehovot, Israel). It has been shown to be beneficial for patients with the exacerbating-remitting form of multiple sclerosis (N. Engl. J. Med. [1987] 317: 408).

SUMMARY OF THE INVENTION

It has recently been observed that patients when treated with copolymer-1 have a lower than expected chance of suffering from visual impairments. This is believed to be of considerable advantage to the general well-being of such patients.

Thus, the present invention relates to the use of copolymer-1 in the manufacture of a medicament for the treatment of visual impairment associated with multiple sclerosis.

In an alternative embodiment the invention relates to the use of copolymer-1 in the manufacture of a medicament for the treatment of optic neuritis.

The present invention includes a method of treating a patient suffering from visual impairment related to multiple sclerosis, comprising administering to said patient a therapeutically effective amount of copolymer-1.

The present invention further includes a method of treating a patient suffering from optic neuritis, comprising administering to said patient a therapeutically effective amount of copolymer
1.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

According to the present invention treatment with copolymer-1 may result in a prevention of any deterioration in visual impairment associated with MS, or in a reduction in the rate of such deterioration. Said treatment may also result in an improvement in vision.

5

10

15

20

25

Examples of the types of visual impairment associated with MS include deteriorations in visual acuity, state of the optic disc, pupil reaction, visual field and ocular motility. Particular conditions that are associated with visual impairment in MS include optic neuritis, retrobulbar optic neuritis, diplopia, dimness of vision and scotomas.

Similarly in the treatment of optic neuritis, treatment with copolymer-1 may result in prevention of further episodes of optic neuritis, delaying of further episodes or return vision to an unimpaired state.

Copolymer-1, according to the present invention, may be prepared by methods known in the art, for example, the process disclosed in US Patent 3,849,550, wherein the N-carboxyanhydrides of tyrosine, alanine, y-benzyl glutamate and E-N-trifluoro-acetyllysine are polymerised at ambient temperature in anhydrous dioxane with diethylamine as initiator. The deblocking of the y-carboxyl group of the glutamic acid is effected by hydrogen bromide in glacial acetic acid and is followed by the removal of the trifluoroacetyl groups from the lysine residues by 1M piperidine. As used herein the terms "ambient temperature" and "room temperature" are used to indicate temperatures from about 20°C to about 26°C.

Compositions of use in the present invention may be formulated by conventional methods known in the art. Preferably, the composition is lyophilized and formed into an aqueous solution suitable for subcutaneous injection, preferably copolymer-1 is formulated with mannitol. Alternatively, copolymer-1 may be formulated in any of the forms known in the art for preparing oral, nasal, buccal, or rectal formulations of peptide drugs.

Typically, copolymer-1 is administered daily to patients at a dosage of 20mg.

The invention will be exemplified but not necessarily limited to the following Examples.

WO 95/33475 PCT/EP95/02125

4

EXAMPLE 1

liters of dioxane.

5

10

15

20

Preparation of Trifluoroacetyl-Copolymer-1

Protected copolymer-1 is prepared as described by Teitelbaum et al. Eur. J. Immun. Vol. 1 p. 242 (1971) from the N-carboxyanhydrides of tyrosine (18g), alanine (50g), γ -benzyl glutamate (35g) and trifluoroacetyllysine (83g) dissolved in 3.5

The polymerization process is initiated by the addition of 0.01 - 0.02% diethylamine. The reaction mixture is stirred at room temperature for 24 hours and then poured into 10 liters water. The product (protected copolymer-1) is filtered, washed with water and dried. The removal of the gamma-benzyl blocking groups from the glutamate residue is carried-out by treating the protected copolymer-1 with 33% hydrobromic acid in glacial acetic acid at room temperature for 6-12 hours with stirring. The product is poured into excess water, filtered, washed and dried, yielding the trifluoroacetyl-copolymer-1.

25 Deprotection of copolymer-1

20g of trifluoroacetyl-copolymer-1 are dispersed in 1 liter of water to which 100g piperidine are added. The mixture is stirred for 24 hours at room temperature and filtered. The solution of crude copolymer-1 is distributed into dialysis bags and dialyzed at 10°-20°C against water until a pH=8 is attained. It is then dialyzed against about 0.3% acetic acid and again water until a pH=5.5-6.0 is obtained. This solution is then concentrated and lyophilized to dryness.

35

30

EXAMPLE 2

Assessment of visual impairment in patients suffering from multiple sclerosis

- 5 Patients were recruited into the study having fulfilled the following criteria;
 - be 18 to 50 years of age,
 - have definite MS as defined by Poser et al. (Ann. Neurol. (1983) 13 227-231),
- be of the relapsing-remitting or relapsing-progressive type when admitted to the trial,
 - have objective evidence of neurological disease that reflects predominantly white matter damage, and
- have had at least two well documented attacks in the two-year period leading up to study entry.

Copolymer-1 was administered sub-cutaneously at a daily dose of 20mg, formulated in 40mg of mannitol.

20 Parameters of visual impairment were measured in the opthalmically acceptable manner as known in the art.

Visual impairment was assessed at six monthly intervals and the total change over the two year period examined.

25

30

35

Results

61 patients completed the full two years of the study. Of these patients, 29 did not experience any relapse of MS during the two years.

Out of the patients experiencing a relapse of MS whilst being treated with copolymer-1, those who had previous experience of optic neuritis (ON) had a 50% chance of having a further ON experience; Patients who had no previous history of ON had a 1 in 14 chance of developing ON. These chances are considerably lower than would normally have been expected had these patients

not been receiving copolymer-1.

Table 1 below shows that in all visual parameters assessed there was a prevention in the deterioration in visual impairment and in many instances there was an improvement as compared to the degree of visual impairment at the commencement of the study. (RE= right eye, LE= left eye)

Table 1

		Improved	No	Worsened	Total tested
Visual acuity	RE	8	43	6	57
	LE	9	44	4	57
Optic Disc	RE	11	38	10	59
	LE	15	40	4	59
Pupil	RE	11	40	8	59
Reaction	LE	7	46	6	59
Visual	RE	6	53	0	59
Field	LE	4	53	2	59
Ocular	RE	2	53	6	61
Motility	LE	2	54	5	61

CLAIMS

- 1. The use of copolymer-1 in the manufacture of a medicament for the treatment of visual impairment associated with multiple sclerosis.
- 5 2. The use of copolymer-1 in the manufacture of a medicament for the treatment of optic neuritis.
 - 3. The use of copolymer-1 in the manufacture of a medicament for the treatment of visual impairment associated with optic neuritis.
- 4. The use of copolymer-1 in the manufacture of a medicament foruse in slowing the deterioration of visual impairment in optic neuritis.
 - 5. A method of treating a patient suffering from visual impairment related to multiple sclerosis, comprising administering to said patient a therapeutically effective amount of copolymer-1.
- A method of treating a patient suffering from optic neuritis,
 comprising administering to said patient a therapeutically effective amount of copolymer-1.
 - 7. The use according to any of Claims 1 to 4 wherein the medicament contains 20mg copolymer-1.
- 8. The method according to any of Claims 5 to 6 comprising daily administration of 20mg copolymer-1.

INTERNATIONAL SEARCH REPORT

Interna: u Application No PCT/EP 95/02125

A. CLASSII	FICATION OF SUBJECT MATTER A61K38/00	
	•	
A crording to	International Patent Classification (IPC) or to both national classification and IPC	
D ETEL DS	SEARCHED	
	nonmentation searched (classification system followed by classification symbols)	1
IPC 6	A61K	
	ion searched other than minimum documentation to the extent that such documents are included in the field	s searched
Documentati	ion service outs. Care	
	·	
Electronic d	ata base consulted during the international search (name of data base and, where practical, search terms us	id)
C DOCIN	IENTS CONSIDERED TO BE RELEVANT	
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
		1 5
X	JOURNAL OF THE NEUROLOGICAL SCIENCES,	1,5
	vol. 31, no. 3, 1977 pages 433-438,	
	l nnën appamski fi Al. 'Effect of a	
'	synthetic polypeptide (COP 1) on patients	
: :	with multiple sclerosis and with acute disseminated encephalomyelitis	
	cited in the application	0.4.6.0
Y	see abstract	2-4,6-8
	see table 1	
x	CLINICAL NEUROPHARMACOLOGY,	1,5
"	vol. 10, no. 5, 1987	
	pages 389-396, LOREN A. ROLAK 'Copolymer-I therapy for	
	multiple sclerosis'	
Υ	see page 391, paragraph 4	2-4,6-8
1	-/	
-	they designed a see listed in the continuation of box C. Palent family members are in	sted in annex.
X Fur	ther documents are listed in the continuation of box C. Patent family members are in	
* Special ca	ategories of cited documents: T later document published after th or priority date and not in confli	et with the application but
'A' docum	nent defining the general state of the art which is not cited to understand the principle dered to be of particular relevance invention.	or theory underlying the
	document but published on or after the international "X" document of particular relevance	unot de comideren w
11. 4000	nent which may throw doubts on priority daim(s) or involve an inventive step when t	r the claimed invention
citatio	on or other special reason (as specified) carnot be considered to involve	an inventive treb when me
other	means ments, such combination being in the art.	DOMORE OF S DELEGIS STATION
P docum	nent published prior to the international filing date but than the priority date claimed & document member of the same	
Date of th	e actual completion of the international search Date of mailing of the internation	nai search report
	4 October 1995 27. 10. 95	
<u></u>	Authorized officer	
Name and	European Patent Office, P.B. 5818 Patentiaan 2	
	NL - 2280 HV Rijewijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl. Tzschoppe. D	•
1	Fax: (+31-70) 340-3016	

INTERNATIONAL SEARCH REPORT

Interns ul Application No PCT/EP 95/02125

		PC1/EP 95/02125
C.(Continue	tion) DOCUMENTS CONSIDERED TO BE RELEVANT	Relevant to daim No.
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Recome to dam not
Y	AM. FAM. PHYS., vol. 48, no. 2, 1993 pages 273-276, MARK J. RIEUMONT ET AL. 'Neuroimaging evaluation in multiple sclerosis' see page 273, right column see page 276, left column	2-4,6-8
Y	PROC. NATL. ACAD. SCI. USA, vol. 91, no. 11, May 1994 pages 4872-4876, MASHA FRIDKIS-HARELI ET AL. 'Direct binding of myelin basic protein and synthetic copolymer 1 to class II major histocompatibility complex molecules on living antigen-presenting cellsspecificity and promiscuity' see abstract	2-4,6-8
Y	ROBERT BERKOW ET AL. 'The Merck Manual of Diagnosis and Therapy' 1992 , MERCK RESEARCH LABORATORIES , RAHWAY N.J. see page 2392 - page 2393	2-4,6-8
<u>-</u>		-
		·
-		

3